arXiv:1506.02421v3 [hep-th] 12 Nov 2015

Irregular matrix model with $\mathcal{W}$ symmetry

Sang Kwan Choi[1] and Chaiho Rim[2]
Department of Physics and Center for Quantum Spacetime (CQUeST)
Sogang University, Seoul 121-742, Korea

Abstract

We present the irregular matrix model which has $\mathcal{W}_3$ and Virasoro symmetry. The irregular matrix model is obtained using the colliding limit of the Toda field theories and produces the inner product between irregular modules of $\mathcal{W}_3$ symmetry. We evaluate the partition function using the flow equation which is the realization of the Virasoro and $\mathcal{W}$-symmetry.

1 Introduction

The irregular matrix model [1] is obtained by the colliding limit of the β-deformed Penner-type matrix model [2, 3] which is equivalent to the regular conformal block of primary fields. The colliding limit [4] is the fusion of primary fields at one point with their Virasoro momenta infinite and results in a non-trivial limit. Especially, there appears irregular module of rank n which is defined as the eigenvector of positive Virasoro generators $L_n, L_{n+1}, \cdots, L_{2n}$. The irregular module of rank 1 can be constructed as the combination of primary and descendant states [5]. However, for the rank greater than 1 the irregular module is not simple to construct because one needs to take account of the consistency condition [6] for the non-negative Virasoro Generators such as $L_0, L_1, \cdots, L_{n-1}$. The consistency condition is not easy to manipulate algebraically. One may detour this difficulty if one uses the irregular matrix model and its conformal symmetry. In our previous papers, we investigate the case with Virasoro symmetry [1, 6, 7, 8]. We may extend this idea to $\mathcal{W}$ symmetry by considering multi-matrix model. $\mathcal{W}$ symmetry was previously used to construct multi-matrix model of polynomial type in [9]. In this paper, we are going to construct the irregular matrix model with $\mathcal{W}$ symmetry using colliding limit of the Toda field theory and present how to find the partition function using the $\mathcal{W}$ symmetry.

Toda field theory is the generalization of the Liouville field theory and contains not only Virasoro symmetry but also higher spin symmetries which is summarized in terms of W-symmetry. Two dimensional Toda field theory associated with simple Lie algebra

---

[1] email:hermit1231@sogang.ac.kr
[2] email:rimpine@sogang.ac.kr

with rank k is defined by the Lagrangian

$$L = \frac{1}{8\pi}(\partial_a \vec{\varphi})^2 + \mu \sum_{i=1}^{k} e^{b\, \vec{e}_i \cdot \vec{\varphi}} \tag{1.1}$$

where $\vec{e}_1, \cdots, \vec{e}_k$ are the simple roots of the Lie algebra. The bosonic vector field $\vec{\varphi}$ has k independent components. The Toda field theory is conformal provided there is a background charge $\vec{Q} = (b + 1/b)\vec{\rho}$ with ρ is the Weyl vector, half of the sum of all positive roots. In this case, the conformal dimension of the exponential terms is 1 and the central charge of the system is $c = k + 12\vec{Q}^2$.

The Toda field theory has k holomorphic symmetry currents $W_2, W_3 \cdots W_{k+1}$ where W_2 is identified as the Virasoro current. In this paper we are concentrated on the colliding limit of the A_2 Toda field theory and construct the A_2 irregular matrix model and find the partition function using the Virasoro and W_3 symmetry. The generalization is straight-forward.

The paper is organized as follows. Section 2 is devoted to the irregular matrix model. Starting with A_2-Toda field theory, we obtain the irregular matrix model using the colliding limit. In section 3, we investigate the $\mathcal{W}$ symmetry in detail. The explicit representation of the $\mathcal{W}$ symmetry is given as the differential operator of the potential variables. In section 4, we present how to evaluate the partition function of the irregular matrix model only using the symmetries. Section 5 is the summary and discussion.

2 A_2 Irregular matrix model

A_2 irregular matrix model is given by two matrix model and is obtained from the colliding limit of the Penner-type two matrix model. For concreteness and comparison, we will use the representation as appeared in [10]. A_2 has two simple roots which will be represented as $\vec{e}_1 = (1, -1, 0)$ and $\vec{e}_2 = (0, 1, -1)$. The simple roots have the squared length 2. The Toda field $\vec{\varphi}$ is conveniently put as $\vec{\varphi} = -\frac{\varphi_1}{\sqrt{6}}(1, 1, -2) - \frac{\varphi_2}{\sqrt{2}}(1, -1, 0)$ whose components are orthogonal to each other. In this convention, the bosonic component satisfies the free field correlation $\varphi_i(z, \bar{z})\varphi_j(w, \bar{w}) \sim -\delta_{ij} \log |z - w|^2$.

The vertex primary field $V_a(z_a) = e^{\vec{\alpha}_a \cdot \vec{\varphi}(z_a)}$ has the conformal dimension $\Delta_a = \vec{\alpha}_a(\vec{Q} - \frac{1}{2}\vec{\alpha}_a)$. We consider the $(n + 2)$-primary field correlation and put the conformal block (holomorphic part of the correlation) in terms of the Selberg integral representation using the screening operator (exponential terms in the Toda Lagrangian). One may put the fields, say, one at the infinity $z_{n+1} \to \infty$, one at the origin $z_0 = 0$, and the rest at z_a $(a = 1, \cdots, n)$. In this case, the conformal block has the form $\left(\prod_{0 \le k < \ell \le n}(z_a - z_b)^{-\vec{\alpha}_a \cdot \vec{\alpha}_b}\right) \times Z_\beta$ if one normalizes the result so that the infinite factor z_{n+1} scales away. The front multiplicative factor is from the free correlation between the primary fields and the rest Z_β is due to the correlation between the screenings and also with the primaries;

$$Z_\beta \equiv \int \prod_{i=1}^{N_1} dx_i \prod_{j=1}^{N_2} dy_j \; \Delta(x)^{2\beta} \Delta(y)^{2\beta} \Delta(x, y)^{-\beta} e^{\frac{2b}{\hbar}\left[\sum_i V_1(x_i) + \sum_j V_2(y_j)\right]} \tag{2.1}$$

where $\Delta(x) = \prod_{i<k}(x_i - x_j)$ and $\Delta(x, y) = \prod_{i,j}(x_i - y_j)$ are the Vandermonde determinant which come from the correlation between screening terms. We put $\beta = -b^2$ to make the partition function similar to the β-deformed matrix model. N_1 and N_2 are number of screening terms with the root vector $\vec{e}_1$ and $\vec{e}_2$ respectively. The number satisfies the neutrality condition [11]

$$\sum_{i=0}^{n} \vec{\alpha}_i + \vec{\alpha}_\infty + b \sum_{k=1}^{2} N_k \vec{e_k} = \vec{Q}\,. \tag{2.2}$$

The potential V_1 and V_2 appear from the correlation between the screening and primary vertex operators. If one parametrizes $\vec{\alpha}_a = \frac{\alpha_a}{\sqrt{3}}(1, 1, -2) + \beta_a(1, -1, 0)$, one has the Penner-type potential

$$\frac{1}{\hbar}V_1(x) = -\sum_{a=0}^{n} \beta_a \log(x - z_a)\,, \quad \frac{1}{\hbar}V_2(y) = -\sum_{a=0}^{n} \frac{1}{2}(\sqrt{3}\alpha_a - \beta_a) \log(y - z_a)\,. \tag{2.3}$$

The irregular matrix model is obtained if one uses the colliding limit similar to the Liouville case [1, 4]: $\alpha_a, \beta_a \to \infty$ and $z_a \to 0$ so that $c_k \equiv \sum_{a=0}^{n} \alpha_a z_a^k$ and $b_k \equiv \sum_{a=0}^{n} \beta_a z_a^k$ with $k = 0, 1, \cdots, n$ are finite. In this case, one has the A_2 irregular matrix model $\mathcal{Z}_n$ with rank n whose potential is given in terms of logarithmic and finite number of inverse powers

$$\frac{1}{\hbar}V_1(z) = -b_0 \log z + \sum_{k=1}^{n} \frac{b_k}{kz^k}\,, \quad \frac{1}{\hbar}V_2(z) = -\tilde{c}_0 \log z + \sum_{k=1}^{n} \frac{\tilde{c}_k}{kz^k}\,. \tag{2.4}$$

where we use the short-hand notation $\tilde{c}_\ell = (\sqrt{3}c_\ell - b_\ell)/2$. In the following, we equally put the irregular matrix model using the new parameter $\hbar \equiv -2ig$ so that

$$\mathcal{Z}_n = \int \prod_{i=1}^{N_1} \prod_{j=1}^{N_2} dx_i dy_j \Delta(x)^{2\beta} \Delta(y)^{2\beta} \Delta(x, y)^{-\beta} e^{-\frac{\sqrt{\beta}}{g}\left[\sum_i V_1(x_i) + \sum_j V_2(y_j)\right]}\,. \tag{2.5}$$

3 $\mathcal{W}$-symmetries of the irregular matrix

3.1 Loop equations

The irregular matrix model (2.5) has two loop-equations. One equation is given as the quadratic equation of the resolvent with its multipoint defined as

$$R_{K_1,\cdots,K_s}(z_1; \cdots; z_s) = \beta \left(\frac{g}{\sqrt{\beta}}\right)^{2-s} \left\langle \sum_{i_1=1}^{N_{K_1}} \frac{1}{z_1 - \lambda_{i_1;K_1}} \cdots \sum_{i_s=1}^{N_{K_s}} \frac{1}{z_s - \lambda_{i_s;K_s}} \right\rangle_{\text{connected}}\,. \tag{3.1}$$

Denoting $\lambda_{i;1} = x_i$, $\lambda_{j;2} = y_j$, one has

$$\begin{aligned}&R_1(z)^2 + R_2(z)^2 - R_1(z)R_2(z) - V_1'(z)R_1(z) - V_2'(z)R_2(z)\\&+ \frac{\hbar Q}{2}\left(R_1'(z) + R_2'(z)\right) - \frac{\hbar^2}{4}\left(R_{1;1}(z,z) - R_{1;2}(z,z) + R_{2;2}(z,z)\right) = \frac{f_1(z)+f_2(z)}{4}\end{aligned} \tag{3.2}$$

where the quantum correction f_1, f_2 are defined as $f_1(z) := 4g\sqrt{\beta}\sum_i^{N_1}\left\langle \frac{-V_1'(z)+V_1'(x_i)}{z-x_i}\right\rangle$ and $f_2(z) := 4g\sqrt{\beta}\sum_i^{N_2}\left\langle \frac{-V_2'(z)+V_2'(y_j)}{z-y_j}\right\rangle$. Here $\langle\cdots\rangle$ denotes the expectation value within the matrix integral. This loop equation is obtained if one performs the conformal transformation of the integration variables $x_i \to x_i + \varepsilon/(x_i - z)$ and $y_j \to y_j + \varepsilon/(y_j - z)$.

The other loop equation is given as a cubic equation [12, 13, 14]:

$$\begin{aligned}0 =& -R_1^2R_2 + R_1R_2^2 - V_1'(R_1^2 + V_1'R_1 - \frac{f_1}{4}) + V_2'(R_2^2 + V_2'R_2 - \frac{f_2}{4}) + \frac{g_1-g_2}{4}\\&+ \frac{\hbar Q}{4}\left[3(V_2'R_2' - V_1'R_1') + R_1R_2' - R_1'R_2 + 2(R_2R_2' - R_1R_1') + V_2''R_2 - V_1''R_1 + \frac{f_1'-f_2'}{4}\right]\\&+ \frac{\hbar^2Q^2}{8}(R_2'' - R_1'') + \frac{\hbar^2}{4}\left[V_1'R_{1;1} - V_2'R_{2;2} + R_{1;1}R_2 - R_{2;2}R_1 - 2R_{1;2}(R_2 - R_1)\right]\\&+ \frac{\hbar^3Q}{16}\left[R_{1;1}' - R_{2;2}' + \lim_{\bar{z}\to z}\left(\frac{\partial}{\partial z}R_{1;2}(z,\bar{z}) - \frac{\partial}{\partial\bar{z}}R_{1;2}(z,\bar{z})\right)\right] + \frac{\hbar^4}{16}(R_{1;2;2} - R_{1;1;2}),\end{aligned} \tag{3.3}$$

where $g_1(z) := 4g^2\beta\sum_{i,j}\left\langle \frac{V_1'(x_i)-V_1'(z)}{(z-x_i)(x_i-y_j)}\right\rangle$ and $g_2(z) := 4g^2\beta\sum_{i,j}\left\langle \frac{V_2'(y_j)-V_2'(z)}{(z-y_j)(y_j-x_i)}\right\rangle$. This is obtained after varying the integration variables $x_i \to x_i + \sum_{j=1}^{N_2}\frac{\epsilon}{(x_i-z)(x_i-y_j)}$ and $y_j \to y_j + \sum_{i=1}^{N_1}\frac{\epsilon}{(y_j-z)(x_i-y_j)}$.

In this paper, we shall focus on the lowest order of $\hbar$ (also putting $Q = 0$). Then, the two equations can be put in a more compact form if we define new notations as $R = R_1 - R_2/2$, $\tilde{R} = \sqrt{3}R_2/2$, $\partial\phi_2 = V_1'$ and $\partial\phi_1 = \frac{1}{\sqrt{3}}(2V_2' + V_1')$.

$$(2R - \partial\phi_2)^2 + (2\tilde{R} - \partial\phi_1)^2 = -\hbar^2\xi_2 \tag{3.4}$$

$$(2\tilde{R} - \partial\phi_1)^3 - 3(2\tilde{R} - \partial\phi_1)(2R - \partial\phi_2)^2 = -\hbar^3\xi_3 \tag{3.5}$$

where ξ_2 and ξ_3 are given explicitly as

$$-\hbar^2\xi_2 = (\partial\phi_2)^2 + (\partial\phi_1)^2 + f_1 + f_2 \tag{3.6}$$

$$\begin{aligned}-\hbar^3\xi_3 =& -(\partial\phi_1)^3 + 3(\partial\phi_2)^2\partial\phi_1 + 3(\partial\phi_1 - \sqrt{3}\partial\phi_2)(f_1 + f_2) + 3\sqrt{3}\partial\phi_2 f_1\\&- \frac{3}{2}(3\partial\phi_1 - \sqrt{3}\partial\phi_2)f_2 - 3\sqrt{3}(g_1 - g_2)\,.\end{aligned} \tag{3.7}$$

ξ_2 and ξ_3 look very complicated but turn out to be the representation of W_2, W_3 currents respectively. This is investigated in the following two subsections.

3.2 Virasoro current

One may use the explicit form of the potential (2.4) to put $\partial\phi_1 = -\hbar\sum_{k=0}^{n} c_k/z^{k+1}$, $\partial\phi_2 = -\hbar\sum_{k=0}^{n} b_k/z^{k+1}$ and

$$f_1(z) + f_2(z) = -\sum_{k=0}^{n-1} \frac{1}{z^{k+2}} \frac{v_k(\hbar^2 \mathcal{Z}_n)}{\mathcal{Z}_n} \tag{3.8}$$

where we use the translation invariance to put $\left\langle \sum_i V_1'(x_i) + \sum_j V_2'(y_j) \right\rangle = 0$. In addition, the Virasoro differential operator is generalized from the one-matrix model, which has the form $v_k = v_k^\partial$ where $v_k^\partial \equiv \sum_{\substack{0\le r\le n\\ r-s=k}} s\left(b_r \frac{\partial}{\partial b_s} + c_r \frac{\partial}{\partial c_s}\right)$. We use the separate notation v_k^∂ for later convenience since v_k will be modified later for the mode $k<0$.

Therefore, ξ_2 in (3.6) has the form

$$\xi_2 = \sum_{k=n}^{2n} \frac{A_k}{z^{k+2}} + \sum_{k=0}^{n-1} \frac{1}{z^{k+2}} \frac{(A_k + v_k^\partial)\mathcal{Z}_n}{\mathcal{Z}_n} \tag{3.9}$$

where A_k is a constant

$$A_k = -\sum_{\ell=0}^{k} (b_\ell b_{k-\ell} + c_\ell c_{k-\ell}) \,. \tag{3.10}$$

We use the convention that $b_\ell = 0 = c_\ell$ when ℓ does not belong to the element set $\{0, 1, \cdots, n\}$. The explicit form ξ_2 is identified with the expectation value of Virasoro current

$$\xi_2 = \frac{\langle \Delta | T(z) | I_n \rangle}{\langle \Delta | I_n \rangle}\,, \qquad T(z) = \sum_{k\in\mathbb{Z}} \frac{L_k}{z^{k+2}}\,, \tag{3.11}$$

if one recalls that the irregular module $|I_n\rangle$ of rank n has the eigenvalue A_k of L_k when $n \le k \le 2n$ and 0 when $k > 2n$. In addition, this identification is consistent with the Virasoro generator representation for the parameter set $\{(b_k, c_k) |\ 0 \le k \le n\}$ as the differential operator $\mathcal{L}_k = A_k + v_k$ which is the right action of the irregular module: $L_k|I_n\rangle := \mathcal{L}_k|I_n\rangle$;

$$\mathcal{L}_k = \begin{cases} 0\,, & 2n < k \\ A_k\,, & n \le k \le 2n \\ A_k + v_k\,, & 0 \le k \le n-1\,. \end{cases} \tag{3.12}$$

3.3 $\mathcal{W}_3$ current

Let us rewrite ξ_3 using the parameters of the potential. First note that

$$g_1(z) - g_2(z) = -\hbar \sum_{k=-2}^{n-2} \frac{\hbar^2 b^2}{z^{k+3}} \left\langle \sum_{i,j} \frac{1}{x_i - y_j} \sum_{r=2}^{n-k} \left(\frac{b_{r+k}}{x_i^r} + \frac{\tilde{c}_{r+k}}{y_j^r} \right) \right\rangle \,. \tag{3.13}$$

To put the expectation values as the derivatives of the partition function, we use two identities: one is

$$4g^2\beta\left\langle\sum_{i,j}\frac{1}{x_i-y_j}\frac{1}{y_j^r}\right\rangle=4g^2\beta\sum_{m=1}^{r}\left\langle\sum_{j,\ell}\frac{1}{y_j^m y_\ell^{r+1-m}}\right\rangle+4g\sqrt{\beta}\left\langle\sum_j\frac{V_2'(y_j)}{y_j^r}\right\rangle \tag{3.14}$$

which is obtained if one changes of integration variable $y_j\to y_j+\varepsilon/y_j^r$ in the partition function $\mathcal{Z}_n$. The other is the same as (3.14) with the exchange of $x_i\leftrightarrow y_j$ and $V_2\to V_1$ due to the obvious symmetry; one may obtain the identity using the integration variable change $x_i\to x_i+\varepsilon/x_i^r$.

Using the above two identities (3.14) and its companion, one finds

$$\begin{aligned}g_1(z)-g_2(z)=\sum_{k=-2}^{n-2}\frac{\hbar}{z^{k+3}}\sum_{r=2}^{n-k}&\Bigg[b_{r+k}\left(\hbar^2b^2\sum_{m=1}^{r-1}\left\langle\sum_{i,k}\frac{1}{x_i^m x_k^{r+1-m}}\right\rangle-2\hbar^2b\sum_{p=0}^{n}\left\langle\sum_i\frac{b_p}{x_i^{r+p}}\right\rangle\right)\\&-\tilde c_{r+k}\left(\hbar^2b^2\sum_{m=1}^{r-1}\left\langle\sum_{j,\ell}\frac{1}{y_j^m y_\ell^{r+1-m}}\right\rangle-2\hbar^2b\sum_{p=0}^{n}\left\langle\sum_j\frac{\tilde c_p}{y_j^{r+p}}\right\rangle\right)\Bigg]\,.\end{aligned} \tag{3.15}$$

Note that in the last term there appear the expectation values of the inverse power of y_j up to $(2n+2)$ and the term higher than n is not directly obtained from the derivative of the rank n partition function. To avoid this difficulty, one may use an extended partition function which is partition function with higher rank, *i.e.*, the partition function with the potential with the parameter c_k, b_k with k up to $2n+2$. In this one has the explicit form of ξ_3

$$\begin{aligned}\xi_3=&\sum_{k=2n}^{3n}\frac{M_k}{z^{k+3}}+\sum_{k=n}^{2n-1}\frac{1}{z^{k+3}}\frac{(M_k+\mu_k)\mathcal{Z}_n}{\mathcal{Z}_n}\\&+\sum_{k=0}^{n-1}\frac{1}{z^{k+3}}\frac{(M_k+\mu_k)\mathcal{Z}_{2n-k}}{\mathcal{Z}_{2n-k}}+\sum_{k=-2}^{-1}\frac{1}{z^{k+3}}\frac{\mu_k(\mathcal{Z}_{2n-k})}{\mathcal{Z}_{2n-k}}\Bigg|_{\{b_{k>n},c_{k>n}\}\to0}\end{aligned} \tag{3.16}$$

where M_k is a constant

$$M_k=\sum_{r+s+t=k}(3b_r\,b_s\,c_t-c_r\,c_s\,c_t)\,. \tag{3.17}$$

The terms with inverse powers of $1/z^k$ is carefully rewritten for $-2\le k\le n-1$ by introducing the extended partition functions, $\mathcal{Z}_{n+1},\cdots,\mathcal{Z}_{2n+2}$. Furthermore, the differential operator μ_k has the different form for the extended case:

$$\mu_k=\begin{cases}\mu_k^{\partial}\,, & n\le k\le 2n-1\\ \mu_k^{\partial}+\mu_k^{\partial^2}\,, & -2\le k\le n-1\end{cases} \tag{3.18}$$

where

$$
\begin{aligned}
\mu_k^{\partial} &\equiv -\sum_{\substack{0\le r,s\le n\\ r+s-t=k}} \frac{t}{2}\left(6b_r c_s \frac{\partial}{\partial b_t} + 3b_r b_s \frac{\partial}{\partial c_t} - 3c_r c_s \frac{\partial}{\partial c_t}\right)\,,\\
\mu_k^{\partial^2} &\equiv \sum_{\substack{0\le r\le n\\ r-s-t=k}} \frac{s\,t}{4}\left(6b_r \frac{\partial}{\partial b_s}\frac{\partial}{\partial c_t} + 3c_r \frac{\partial}{\partial b_s}\frac{\partial}{\partial b_t} - 3c_r \frac{\partial}{\partial c_s}\frac{\partial}{\partial c_t}\right)\,.
\end{aligned} \tag{3.19}
$$

We identify ξ_3 with the expectation value of the $\mathcal{W}_3$ current between a regular module $|\Delta\rangle$ and an irregular module $|I_n\rangle$ as

$$
\xi_3 = \frac{\langle\Delta|W(z)|I_n\rangle}{\langle\Delta|I_n\rangle}\,,\quad W(z) = \sum_{k\in\mathbb{Z}} \frac{W_k}{z^{k+3}}\,. \tag{3.20}
$$

Note that the irregular module has the eigenvalue M_k of W_k when $2n \le k \le 3n$. Higher mode W_k with $k > 3n$ annihilates the irregular module. In this identification one has the representation of the $\mathcal{W}_3$ current $W_k|I_n\rangle := \omega_k|I_n\rangle$ where

$$
\omega_k = \begin{cases} 0\,, & 3n < k\\ M_k\,, & 2n \le k \le 3n\\ M_k + \mu_k\,, & n \le k \le 2n-1\\ M_k + \mu_k^{\partial} + \mu_k^{\partial^2}\,, & 0 \le k \le n-1\,. \end{cases} \tag{3.21}
$$

Considering a negative mode acts on the regular module and vanishes *i.e.*, $\langle\Delta|\omega_k|I_n\rangle = 0$ for $k < 0$, one expects the mode with $k = -1, -2$ in (3.16) to vanish. This is confirmed in the next subsection.

3.4 Consistency check of the representation

One can check the representation (3.12) and (3.21) is compatible with the commutation relation of the Virasoro and $\mathcal{W}_3$ modes[15]:

$$
[L_p, L_q] = (p-q)L_{p+q} + \frac{c}{12}(p^3 - p)\delta_{p,-q}\,, \tag{3.22}
$$

$$
[L_p, W_q] = (2p-q)W_{p+q}\,, \tag{3.23}
$$

$$
\begin{aligned}
-\frac{2}{9}[W_p, W_q] =& \frac{c}{3\cdot 5!}(p^2-1)(p^2-4)p\delta_{p,-q} + \frac{16}{22+5c}(p-q)\Lambda_{p+q}\\
&+ (p-q)\left(\frac{1}{15}(p+q+2)(p+q+3) - \frac{1}{6}(p+2)(q+2)\right)L_{p+q}
\end{aligned} \tag{3.24}
$$

where[1]

$$
\Lambda_p = \sum_{k=-\infty}^{\infty} :L_k L_{p-k}: + \frac{1}{5}x_p L_p\,,
$$

$$
x_{2\ell} = (\ell+1)(\ell-1) \qquad x_{2\ell+1} = (2+\ell)(1-\ell)
$$

---

[1] If one rescales W_p as $i\frac{3}{\sqrt{2}}W_p$, then the algebra reduces to the original Fateev and Zamolodchikov convention [15].

and the central charge $c = 2 + 12\vec{Q}^2$.

Note that the irregular module $|I_n\rangle$ with rank n is defined as a simultaneous eigenstate of L_k's and W_k's:

$$L_k|I_n\rangle = A_k|I_n\rangle\,, \qquad n \le k \le 2n \tag{3.25}$$

$$W_k|I_n\rangle = M_k|I_n\rangle\,, \quad 2n \le k \le 3n \tag{3.26}$$

where A_k and M_k are eigenvalues. In addition, one requires that the action of $L_{k>2n}$ and $W_{k>3n}$ vanish.

The eigenvalues are not enough to define the irregular module. One needs $L_{k<n}$ and $W_{k<2n}$. The Virasoro generator has the representation $\mathcal{L}_k$ for $0 \le k < n$ in (3.12) and $\mathcal{W}_3$ generator ω_k for $-2 \le k < 2n$ in (3.21). However, this representation is not enough to check the commutation relation (3.24). This is because Λ_p contains the negative mode $L_{k<0}$ of Virasoro generators and hence, the non-negative modes are not closed by themselves in the commutation relation. On the other hand, one cannot obtain the information of the negative modes from the equation (3.20) because the negative mode contribution vanishes in the expectation value. Therefore we need another way to find the negative mode representation.

To find the negative modes we consider the identity (3.14) and its companion for the extended partition function $\mathcal{Z}_{n-k}$ for $k < -1$:

$$\left(v_k^{\partial} + v_k^{\partial^2}\right)\mathcal{Z}_{n-k} = 0\,, \qquad v_k^{\partial^2} \equiv -\sum_{-(r+s)=k} \frac{r\,s}{4}\left(\frac{\partial}{\partial b_r}\frac{\partial}{\partial b_s} + \frac{\partial}{\partial c_r}\frac{\partial}{\partial c_s}\right)\,. \tag{3.27}$$

The invariant property of the partition function shows that one has L_k with $v_k = v_k^{\partial} + v_k^{\partial^2}$ for $k < -1$ with the extended set of parameters $\{b_{k\ge n}, c_{k\ge 0}\}$ when necessary.

Likewise for $\mathcal{W}_3$, one has the desired identity if one considers the transformation $x_i \to x_i + \sum_j^{N_2} \frac{\epsilon}{(x_i-y_j)x_i^r}$ and $y_j \to y_j + \sum_i^{N_1} \frac{\epsilon}{(x_i-y_j)y_j^r}$ to get

$$\left(\mu_k^{\partial} + \mu_k^{\partial^2}\right)\mathcal{Z}_{2n-k} = 0 \quad \text{for } k = -1, -2 \tag{3.28}$$

$$\left(\mu_k^{\partial} + \mu_k^{\partial^2} + \mu_k^{\partial^3}\right)\mathcal{Z}_{2n-k} = 0 \quad \text{for } k \le -3\,, \tag{3.29}$$

where

$$\mu_k^{\partial^3} \equiv -\sum_{-(r+s+t)=k} \frac{r\,s\,t}{8}\left(3\frac{\partial}{\partial b_r}\frac{\partial}{\partial b_s}\frac{\partial}{\partial b_t} - \frac{\partial}{\partial c_r}\frac{\partial}{\partial c_s}\frac{\partial}{\partial c_t}\right)\,. \tag{3.30}$$

Eq (3.28) shows that one has $\mu_k(\mathcal{Z}_{2n-k}) = 0$ for $k = -1, -2$, or $\langle\Delta|\omega_k|I_n\rangle = 0$ as asserted in sec 3.3. The invariant property of the partition function (3.29) allows to put $\mu = \mu_k^{\partial} + \mu_k^{\partial^2} + \mu_k^{\partial^3}$ for $k \le -3$.

In fact, the negative mode representation is easily understood if one notes that this representation is found from the coherent state representation of Heisenberg algebra whose positive mode a_k has the eigenvalues b_k or c_k when $k > 0$. In the coherent state representation, the negative mode is given as the differential operator $b_{-k} = -(k/2)\partial/(\partial b_k)$ when $k > 0$. This is the reason why one and two-derivative terms appear in the negative Virasoro mode representation while one, two and three-derivative terms appear in the negative W mode representation as appeared in [9].

4 Irregular partition function

4.1 Differential equations

The loop equations (3.4) and (3.5) give a series of differential equations for the partition function $\mathcal{Z}_n$. Large z expansion gives $2n$-differential equations:

$$-v_k \log \mathcal{Z}_n = d_k \ , \quad 0 \le k \le n-1 \tag{4.1}$$

$$-\mu_k \log \mathcal{Z}_n = e_k \ , \quad n \le k \le 2n-1 \tag{4.2}$$

where v_k and μ_k are differential operators defined in (3.12), (3.21). One may also find differential equations corresponding to μ_k for $k < n$. However, this equation contains the extended set of parameters $\{b_{k>n}, c_{k>n}\}$ and is redundant since we have $2n$-equations (4.1), (4.2) which will completely fix the partition function. d_k and e_k in (4.1) and (4.2) are given as

$$d_k = A_k + \sum_{r+s=k} \left(d^b_r\, d^b_s + d^c_r\, d^c_s \right) \ , \quad e_k = M_k + \sum_{r+t+s=k} \left(d^c_r\, d^c_t\, d^c_s - 3 d^b_r\, d^b_t\, d^c_s \right) \tag{4.3}$$

$$d^c_r = \frac{\sqrt{3}}{2} \left\langle \sum_j y_j^r \right\rangle + c_r \ , \qquad d^b_r = \left\langle \sum_i x_i^r \right\rangle - \frac{1}{2} \left\langle \sum_j y_j^r \right\rangle + b_r \, .$$

Here we use the convention of the coefficients $b_k = c_k = 0$ when $k > n$. Note that d_k and e_k in (4.3) are unknown except d_0 since they are given in terms of expectation values. One has to find these expectation values as the function of the coefficients of the potential explicitly. This can be done by finding the filling fraction with the resolvents $R_1(z)$ and $R_2(z)$ across a given branch cut. This idea is used to evaluate the partition function in section 4.3.

4.2 Loop equations and spectral curve

The resolvents R_1 and R_2 have the asymmetric role in the loop equation. This is because we put the loop equation asymmetrically. One may put the loop equation in a symmetric way. Suppose we introduce

$$u_1(z) := R_1(z) + t_1(z) \, , \qquad t_1(z) = -\frac{2V_1'(z) + V_2'(z)}{3} \, , \tag{4.4}$$

$$u_2(z) := -R_2(z) + t_2(z) \, , \qquad t_2(z) = \frac{V_1'(z) + 2V_2'(z)}{3} \tag{4.5}$$

and $u_0(z) := -u_1(z) - u_2(z)$. Then, using the two loop equations (3.4) and (3.5) one finds a spectral curve in a cubic form [12, 13, 14, 16]

$$\prod_{i=0}^{2} (u - u_i(z)) = u^3 + \frac{\hbar^2 \xi_2(z)}{4} u - \frac{\hbar^3 \xi_3(z)}{12\sqrt{3}} = 0 \, . \tag{4.6}$$

This shows that u_2 is the solution of the cubic equation which is exactly the same form as the loop equation given in (3.4) and (3.5). The spectral curve also demonstrates that



Figure 1: Covering space and its cut structure

u_1 and u_0 are two other solutions. Therefore, u_1 and u_0 should respect the same loop equation (3.4) and (3.5).

To understand the three branches u_0, u_1, u_2, we first the case with no quantum correction, *i.e.*, $f_1 = f_2 = g_1 = g_2 \equiv 0$. In this case ξ_2 and ξ_3 are given in terms of the potentials only and the three roots u_1, u_2, u_0 are reduced to t_1, t_2 and $t_0(z) := -t_1(z) - t_2(z)$ defined in (4.4), (4.5). The classical spectral curve has common points. For example, when $t_0(z) = t_1(z)$ one finds $V_1'(z) = 0$. The stationary point of the potential V_1 corresponds to the double points, the intersect of the branches t_0 and t_1. Likewise, $t_0(z) = t_2(z)$ corresponds to the case $V_2'(z) = 0$, and $t_1(z) = t_2(z)$ to $V_1'(z) + V_2'(z) = 0$. This shows that the three classical branches are connected to each other at the stationary points of the potentials.

If the spectral curve is deformed by f_1, f_2, g_1 and g_2, then the double point splits and forms a branch cut. As a result, $3n$-number of branch cuts appear on the complex plane z when all the double points are distinct. Let us denote the branch cuts as $I^{(i)}_{[ab]}$ connecting two branches u_a and u_b with $i = 1, \cdots, n$ (see Fig.1).

One may count the number of eigenvalues of the potential (number of integration variables) by taking the contour integral of the resolvent around the cut. Let us denote the contour around the cut $I^{(i)}_{[k,k+1]}$ as $\mathcal{A}^{(i)}_{[k,k+1]}$ assuming the branch index $k \bmod 3$. One may find the filling fractions (number of eigenvalues) using u_k,

$$\frac{\hbar b}{2} n^{(i)}_{k[k+1]} := \frac{1}{2\pi i} \oint_{\mathcal{A}^{(i)}_{[k,k+1]}} u_k(z) dz \tag{4.7}$$

or using u_{k+1},

$$\frac{\hbar b}{2} n^{(i)}_{k+1[k]} := \frac{1}{2\pi i} \oint_{\mathcal{A}^{(i)}_{[k,k+1]}} u_{k+1}(z) dz \,. \tag{4.8}$$

These two quantities add up to zero; $n^{(i)}_{k[k+1]} + n^{(i)}_{k+1[k]} = 0$ since

$$u_k(\lambda + i0) - u_k(\lambda - i0) = -\big(u_{k+1}(\lambda + i0) - u_{k+1}(\lambda - i0)\big) \quad \text{for} \quad \lambda \in I^{(i)}_{[k,k+1]} \,. \tag{4.9}$$

This is due to the fact that $u_k + u_{k+1} + u_{k+2} = 0$ and u_{k+2} is continuous on the cut $I^{(i)}_{[k,k+1]}$. Therefore, one may have $N = N_1 + N_2$ where

$$N_1 = \sum_{i=1}^{n} \left(n^{(i)}_{1[0]} + n^{(i)}_{1[2]} \right) \,, \quad N_2 = -\sum_{i=1}^{n} \left(n^{(i)}_{2[1]} + n^{(i)}_{2[0]} \right) \,. \tag{4.10}$$

(Note the minus sign in N_2 comes from the definition of u_2 in (4.5).)

4.3 Evaluation of partition function

We evaluate the partition function of the rank 1 explicitly in this subsection. The rank 1 partition function has two differential equations

$$-v_0 \log \mathcal{Z}_1 = d_0 \,, \qquad -\mu_1 \log \mathcal{Z}_1 = e_1 \tag{4.11}$$

where $v_0 = b_1 \frac{\partial}{\partial b_1} + c_1 \frac{\partial}{\partial c_1}$, $\mu_1 = -3b_1 c_1 \frac{\partial}{\partial b_1} - \frac{3}{2}(b_1^2 - c_1^2)\frac{\partial}{\partial c_1}$ and d_0, e_1 are defined in (4.3). Since d_0 is a constant

$$d_0 = (bN_1)^2 - bN_1\, bN_2 + (bN_2)^2 + 2b_0\, bN_1 - b_0\, bN_2 + \sqrt{3} c_0\, bN_2 \tag{4.12}$$

one has the solution of the first equation in (4.11) of the form,

$$\log \mathcal{Z}_1 = -d_0 \log c_1 + H(t) \tag{4.13}$$

where $t := b_1/c_1$ and $H(t)$ is a homogeneous solution to v_0.

Plugging this into the second one in (4.11), one obtains

$$(3 - t^2) t \frac{\partial H(t)}{\partial t} = \frac{2}{3} \frac{e_1}{c_1} + (t^2 - 1) d_0 \,. \tag{4.14}$$

where e_1 is given in terms of the expectation values $\langle \sum_i x_i \rangle$ and $\langle \sum_j y_j \rangle$,

$$\begin{aligned} e_1 =& 3b_0^2 c_1 - 3c_0^2 c_1 + 6b_0 c_0 b_1 + \frac{3}{8}\left(2c_0 + \sqrt{3} N_2\right)^2 \left(2c_1 + \sqrt{3} \langle \sum_j y_j \rangle\right) \\ & - \frac{3}{8}(2b_0 + 2N_1 - N_2) \left[2\Big(2b_1 + 2\langle \sum_i x_i \rangle - \langle \sum_j y_j \rangle\Big) \left(2c_0 + \sqrt{3} N_2\right) \right. \\ & \left. + (2b_0 + 2N_1 - N_2)\left(2c_1 + \sqrt{3}\langle \sum_j y_j \rangle\right) \right] . \end{aligned} \tag{4.15}$$

To solve the equation (4.14), we need the explicit from of e_1/c_1 as the function of t. This can be done using the constraint of the filling fraction.

For the rank 1, we have three cuts $I_{[01]}$, $I_{[12]}$ and $I_{[20]}$ whose classical double points are $-\frac{b_1}{b_0}$, $\frac{-b_1-\sqrt{3}c_1}{b_0+\sqrt{3}c_0}$ and $\frac{-b_1+\sqrt{3}c_1}{b_0-\sqrt{3}c_0}$, respectively. However, one filling fraction is enough, say

$$\frac{1}{2\pi i}\oint_{\mathcal{A}_{[01]}} u_1(z)dz = \frac{\hbar b}{2} n_{1[0]}\,. \tag{4.16}$$

The branch u_1 in (4.6) is given as

$$\frac{1}{\hbar}u_1(z) = \frac{\left(\sqrt{3}+i3\right)\mathcal{P}_2 + \left(\sqrt{3}-i3\right)\left(\mathcal{P}_3+\sqrt{\mathcal{P}_3^2-\mathcal{P}_2^3}\right)^{2/3}}{12\left(\mathcal{P}_3+\sqrt{\mathcal{P}_3^2-\mathcal{P}_2^3}\right)^{1/3}} \tag{4.17}$$

where $\mathcal{P}_2$ and $\mathcal{P}_3$ are polynomials given by

$$\begin{aligned}\mathcal{P}_2(z) &= (b_0^2+c_0^2+d_0)z^2 + 2(b_0b_1+c_0c_1)z + b_1^2 + c_1^2\,,\\ \mathcal{P}_3(z) &= (c_0^3 - 3b_0^2c_0 + e_0)z^3 + (3c_0^2c_1 - 3b_0^2c_1 - 6b_0c_0b_1 + e_1)z^2\\ &\quad - 3(c_0b_1^2 + 2b_0b_1c_1 - c_0c_1^2)z + c_1^3 - 3b_1^2c_1\,.\end{aligned}$$

Here e_0 is a constant, $e_0 = 3b_0^2c_0 - c_0^3 - 3(N_1 - \frac{N_2}{2} + b_0)^2(\frac{\sqrt{3}}{2}N_2 + c_0) + (\frac{\sqrt{3}}{2}N_2 + c_0)^3$. Six roots of $(\mathcal{P}_3^2 - \mathcal{P}_2^3)$ implies three branch cuts on z-plane. (Note that a cubic branch cut whose branch points are roots of $\left(\mathcal{P}_3 + \sqrt{\mathcal{P}_3^2-\mathcal{P}_2^3}\right)$ is not reduced to the classical point and no eigenvalues lie on it. Thus we don't need to take account of this cut.)

To find the filling fraction perturbatively, we assume that $|t| \ll 1$. In this case, the cut $I_{[01]}$ is widely separated from the others: Rescaling the integration variable in the contour integral (4.16) with b_1, the cut $I_{[01]}$ has a finite width whereas other cuts shrink to a point as $t \to 0$. Thus, one can expand the $u_1(z)$ in (4.17) safely in powers of t to get $\left(\mathcal{P}_2^3 - \mathcal{P}_3^2\right) = b_1^2c_1^4\left[Q_0 + tQ_1 + \mathcal{O}(t^2)\right]$. However, the explicit form of Q_i depends on the relative scales of e_1. Small t-expansion assumes that $|b_1| < |c_1|$. In addition, since e_1 is a quantum deformation, $|c_1| < |e_1|$ is not allowed. Therefore, we have two different relative scales: (I) $|e_1| \lesssim |b_1| \ll |c_1|$ and (II) $|b_1| \ll |e_1| \lesssim |c_1|$.

Let us first consider the case (I). In this case $\tilde{e}_1 := e_1/b_1$ is small. Assuming $\tilde{e}_1 = \mathcal{O}(t^0)$ at most, one has

$$\begin{aligned}Q_0(z) &= 3\left((3b_0^2+d_0)z^2 + 6b_0z + 3\right)\\ Q_1(z) &= z\left((36b_0^2c_0 + 12c_0d_0 - 2e_0)z^2 + (72b_0c_0 - 2\tilde{e}_1)z + 36c_0\right)\,.\end{aligned} \tag{4.18}$$

The filling fraction $n_{1[0]}$ is given in power of t,

$$\begin{aligned}bn_{1[0]} &= \frac{1}{\pi i}\oint_{\mathcal{A}_{[01]}} dz\left[\frac{c_0z + \sqrt{Q_0(z)/3}}{2\sqrt{3}z^2} - t\frac{\tilde{e}_1 + e_0z - d_0\sqrt{Q_0(z)/3}}{6\sqrt{3}\sqrt{Q_0(z)/3}} + \mathcal{O}(t^2)\right]\\ &= -b_0 + \sqrt{b_0^2 + \frac{d_0}{3}} - t\frac{(d_0+3b_0^2)\tilde{e}_1 - 3b_0e_0}{3\sqrt{3}(d_0+3b_0^2)^{3/2}} + \mathcal{O}(t^2)\,.\end{aligned} \tag{4.19}$$

Noting $\tilde{e}_1 = \mathcal{O}(t^0)$, one has to require the filling fraction $bn_{1[0]} = -b_0 + \sqrt{b_0^2 + d_0/3} + tb\Delta n_{1[0]}$ so that $\Delta n_{1[0]} = \mathcal{O}(t^0)$. Since we regard $n_{1[0]}$ as the controlling parameter, we force $\Delta n_{1[0]} = 0$ so that

$$bn_{1[0]} = -b_0 + \sqrt{b_0^2 + d_0/3}\,. \tag{4.20}$$

In this case, one has

$$\tilde{e}_1 = \frac{3b_0 e_0}{3b_0^2 + d_0} - t\frac{3\left(d_0(d_0^3 + 2c_0 d_0 e_0 - e_0^2) + b_0^2(3d_0^3 + 6c_0 d_0 e_0 + 2e_0^2)\right)}{4(3b_0^2 + d_0)^3} + \mathcal{O}(t^2) \tag{4.21}$$

and finds the partition function from (4.14)

$$\mathcal{Z}_1 = \mathcal{N}\, b_1^{-\frac{d_0}{3}}\, c_1^{-\frac{2}{3}d_0}\; e^{\frac{2b_0 e_0}{9b_0^2 + 3d_0}t + \mathcal{O}(t^2)} \tag{4.22}$$

where $\mathcal{N}$ is the normalization independent of t.

For the case (II), one finds a quite different structure from that of (I). Note that $|b_1| < |e_1|$. This parameter domain allows to put $b_1 = 0$. In addition, b_1 should be zero when the quantum correction e_1 is zero. In this case, the double point of the classical branches of t_0 and t_1 does not exist on the finite domain of the complex plane since it is now given by the stationary point of V_1, satisfying $V_1'(z) = \frac{b_0}{z} = 0$. Considering this, one may force $n_{1[0]} = 0$ which is true at the classical level. With b_1 which can be 0, we had better introduce a new parameter $\hat{e}_1 := e_1/c_1$ instead of $\tilde{e}_1$. Assuming $\hat{e}_1 = \mathcal{O}(t^0)$ at most, one has

$$\begin{aligned} \mathcal{Q}_0(z) &= (9b_0^2 + 3d_0 - 2\hat{e}_1)z^2 + 18b_0 z + 9 \\ \mathcal{Q}_1(z) &= z\left((36b_0^2 c_0 + 12c_0 d_0 - 2e_0 - 6c_0\hat{e}_1)z^2 + 72b_0 c_0 z + 36c_0\right) \end{aligned} \tag{4.23}$$

and the filling fraction has the form

$$\begin{aligned} bn_{1[0]} &= \frac{1}{\pi i}\oint_{\mathcal{A}_{[01]}} dz \left[\frac{\sqrt{3}c_0 z + \sqrt{\mathcal{Q}_0(z)}}{6z^2} + t\frac{9(c_0\hat{e}_1 - e_0)z + \sqrt{3}(3d_0 + \hat{e}_1)\sqrt{\mathcal{Q}_0(z)}}{54\sqrt{\mathcal{Q}_0(z)}} + \mathcal{O}(t^2)\right] \\ &= -b_0 + \frac{\sqrt{9b_0^2 + 3d_0 - 2\hat{e}_1}}{3} + t\frac{b_0(e_0 - c_0\hat{e}_1)}{(9b_0^2 + 3d_0 - 2\hat{e}_1)^{3/2}} + \mathcal{O}(t^2)\,. \end{aligned} \tag{4.24}$$

Putting $n_{1[0]} = 0$, one finds

$$\hat{e}_1 = \frac{3}{2}d_0 + t\frac{2e_0 - 3c_0 d_0}{2b_0} + \mathcal{O}(t^2) \tag{4.25}$$

and the partition function from (4.14),

$$\mathcal{Z}_1 = \hat{\mathcal{N}} c_1^{-d_0}\, e^{\frac{2e_0 - 3c_0 d_0}{9b_0}t + \mathcal{O}(t^2)} \tag{4.26}$$

with $\hat{\mathcal{N}}$ a normalization independent of t. In this evaluation, we assume that $b_1 > 0$ and the vanishing filling fraction is obtained when $b_0 > 0$. The same is true if one assume

$b_1 < 0$ with $b_0 < 0$. The vanishing filling fraction condition holds as far as $b_0 b_1 > 0$, which is the case V_1 having no stationary point in our complex domain. (Note that due to the logarithmic potential, we exclude the negative real axis in our complex domain).

Suppose we put $b_1 \equiv 0$ from the beginning and require the filling fraction $n_{1[0]}$ to vanish. In this case, one finds that the partition function is simply reduced to $\mathcal{Z}_1 = \hat{\mathcal{N}} c_1^{-d_0}$ since $t = 0$. In addition, the two differential operators v_0 and μ_1 are not independent but has the relation $\mu_1 = \frac{3}{2} c_1 v_0$. The case with $b_1 = 0$ is identified in [10] with the semi-degenerate where the regular module has the null vector at the first level [15, 17, 18]. In our approach, if the semi-degenerate module is put at infinity, we have the semi-degenerate $\langle \Delta |$ in (3.20) whose conformal dimension is $\Delta = \vec{\alpha}_\infty (\vec{Q} - \frac{1}{2}\vec{\alpha}_\infty)$ with $\vec{\alpha}_\infty = \varkappa \vec{w}_2$ where $\vec{w}_2$ is the fundamental weight satisfies $\vec{w}_i \cdot \vec{e}_j = \delta_{ij}$. In this case, we have $N_1 = 0$ as the semi-degenerate result.

5 Summary and discussion

We generalize the (Virasoro) irregular matrix model so that the model contains the Virasoro and $\mathcal{W}_3$ symmetry. This model is constructed using the colliding limit of A_2 Toda field theory. The symmetry of the irregular models is analyzed through the loop equations which have the quadratic and cubic form. The spectral curve obtained corresponds to the Seiberg-Witten curve of the $SU(3)$ super-conformal linear quiver theory.

It should be emphasized the spectral curve (4.6) is enough to find the partition function of the irregular matrix model without evaluation of the functional integral or Selberg integral. Using the differential representation of the Virasoro and $\mathcal{W}_3$ symmetry generators we derive the differential equations for the partition function from the loop equations. The differential equations allow us to find the partition function of the irregular model. We present the explicit form of the representation and find the partition function to the lowest order of $\hbar$ (corresponding to the large N limit) for the non-trivial case (irregular module with rank 1). It is not difficult to find the partition function with rank greater than 1 if one uses the parameter scale as $\left|\frac{b_{k+1}}{b_k}\right| \ll \left|\frac{b_k}{b_{k-1}}\right|$, $\left|\frac{c_{k+1}}{c_k}\right| \ll \left|\frac{c_k}{c_{k-1}}\right|$ and $\left|\frac{b_1}{b_0}\right| \ll \left|\frac{c_n}{c_{n-1}}\right|$ as was used in Liouville case [1, 6].

Once the partition function is known, one may construct the irregular conformal block(ICB), noting that the partition function with appropriate potential is related with an inner product between an irregular module and regular/irregular modules. The simplest ICB, the inner product $\langle I_m | I_n \rangle$ is given in terms of irregular matrix model. One may consider the irregular partition function $\mathcal{Z}_{(m:n)}$ where irregular module of rank

n lies at the origin and one with rank m at infinity:

$$\mathcal{Z}_{(m:n)} = \int \prod_{i=1}^{N_1} \prod_{j=1}^{N_2} dx_i dy_j \Delta(x)^{2\beta} \Delta(y)^{2\beta} \Delta(x,y)^{-\beta} e^{-\frac{\sqrt{\beta}}{g}\left[\sum_i V_1^{(m:n)}(x_i) + \sum_j V_2^{(m:n)}(y_j)\right]} , \tag{5.1}$$

$$\frac{V_1(z)}{\hbar} = -b_0 \log z + \sum_{k=1}^{n} \frac{b_k}{k z^k} + \sum_{\ell=1}^{m} \frac{b_{-\ell} z^\ell}{\ell} , \quad \frac{V_2(z)}{\hbar} = -\tilde{c}_0 \log z + \sum_{k=1}^{n} \frac{\tilde{c}_k}{k z^k} + \sum_{\ell=1}^{m} \frac{\tilde{c}_{-\ell} z^\ell}{\ell} .$$

There are a few subtle issues when one identifies this partition function with the inner product $\langle I_m | I_n \rangle$. One is the extra contribution due to the colliding limit: $\prod_{a,b}(1 - z_b/\bar{z}_a)^{-\vec{\alpha}_a \cdot \vec{\alpha}_b} \to e^{\zeta_{(m:n)}}$ with $\zeta_{(m:n)} = \sum_k^{min(m,n)} 2(b_k b_{-k} + c_k c_{-k})/k$ as $z_b \to 0$, $\bar{z}_a \to \infty$. This non-trivial contribution was considered in [7] for A_1 case. The other is about the normalization for the case with rank greater 1. One has to take account the normalization properly because of the consistency condition for the Virasoro and $\mathcal{W}$ symmetry [6]. This consideration leads to the irregular conformal block $\mathcal{F}^{(m:n)}$;

$$\mathcal{F}^{(m:n)} = \frac{e^{\zeta_{(m:n)}} \mathcal{Z}_{(m:n)}}{\mathcal{Z}_{(0:n)} \mathcal{Z}_{(m:0)}} . \tag{5.2}$$

One may find the complete representation of the symmetry generators which has non-leading order of $\hbar$ with $Q \neq 0$ without difficulty considering the full equations given in (3.2), (3.3). In addition, the irregular matrix model is easy to generalize into A_k model since it is composed of k-matrix potential along with the Vandermonde determinant whose power is given in terms of Dynkin index. There will appear more than cubic power in the loop equations. To control these complicated equations, one may resort to DDAHA as demonstrated in [19]. This will be presented elsewhere in the future.

Finally, the irregular matrix model is motivated by Argyres-Douglas theory [20, 21] in connection with AGT conjecture [22], which develops irregular punctures to the holomorphic one form of the Hitchin system [23, 24, 25]. So far, the Virasoro case has been intensively investigated using the irregular matrix model. Extension to $\mathcal{W}$ symmetry will provide a useful tool to study Argyres-Douglas theory corresponding to $SU(N)$ gauge theory.

Acknowledgement

This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIP) (NRF-2014R1A2A2A01004951).

References

[1] T. Nishinaka and C. Rim, *Matrix models for irregular conformal blocks and Argyres-Douglas theories,* JHEP **1210**, 138 (2012) [arXiv:1207.4480 [hep-th]].

[2] R. Dijkgraaf and C. Vafa, *Toda Theories, Matrix Models, Topological Strings, and N=2 Gauge Systems,* arXiv:0909.2453 [hep-th].

[3] T. Eguchi and K. Maruyoshi, *Penner Type Matrix Model and Seiberg-Witten Theory,* JHEP **1002** (2010) 022 [arXiv:0911.4797].

[4] D. Gaiotto and J. Teschner, *Irregular singularities in Liouville theory,* JHEP **1212** (2012) 050 [arXiv:1203.1052].

[5] D. Gaiotto, *Asymptotically free N=2 theories and irregular conformal blocks* [arXiv:0908.0307].

[6] S.-K. Choi and C. Rim, *Parametric dependence of irregular conformal block,* JHEP **1404** (2014) 106 [arXiv:1312.5535].

[7] S. K. Choi, C. Rim and H. Zhang, *Virasoro irregular conformal block and beta deformed random matrix model,* Phys. Lett. B **742**, 50 (2015) [arXiv:1411.4453 [hep-th]].

[8] C. Rim and H. Zhang, *Classical Virasoro irregular conformal block,* arXiv:1504.07910 [hep-th].

[9] S. Kharchev, A. Marshakov, A. Mironov, A. Morozov and S. Pakuliak, *Conformal matrix models as an alternative to conventional multimatrix models,* Nucl. Phys. B **404**, 717 (1993) [hep-th/9208044].

[10] H. Kanno, K. Maruyoshi, S. Shiba, and M. Taki, W_3 *irregular states and isolated N=2 superconformal field theories* [arXiv:1301.0721].

[11] V. A. Fateev and A. V. Litvinov, *Correlation functions in conformal Toda field theory. I.,* JHEP **0711**, 002 (2007) [arXiv:0709.3806 [hep-th]].

[12] A. Klemm, K. Landsteiner, C. I. Lazaroiu and I. Runkel, *Constructing gauge theory geometries from matrix models,* JHEP **0305**, 066 (2003) [hep-th/0303032].

[13] S. G. Naculich, H. J. Schnitzer and N. Wyllard, *Cubic curves from matrix models and generalized Konishi anomalies,* JHEP **0308**, 021 (2003) [hep-th/0303268].

[14] R. Schiappa and N. Wyllard, *An A(r) threesome: Matrix models, 2d CFTs and 4d N=2 gauge theories,* J. Math. Phys. **51**, 082304 (2010) [arXiv:0911.5337 [hep-th]].

[15] V. A. Fateev and A. B. Zamolodchikov, *Conformal Quantum Field Theory Models in Two-Dimensions Having Z(3) Symmetry,* Nucl. Phys. B **280**, 644 (1987).

[16] R. Dijkgraaf and C. Vafa, *On geometry and matrix models,* Nucl. Phys. B **644**, 21 (2002) [hep-th/0207106].

[17] N. Wyllard, *A(N-1) conformal Toda field theory correlation functions from conformal N = 2 SU(N) quiver gauge theories,* JHEP **0911**, 002 (2009) [arXiv:0907.2189 [hep-th]].

[18] S. Kanno, Y. Matsuo, S. Shiba and Y. Tachikawa, *N=2 gauge theories and degenerate fields of Toda theory,* Phys. Rev. D **81**, 046004 (2010) [arXiv:0911.4787 [hep-th]].

[19] Y. Matsuo, C. Rim and H. Zhang, *Construction of Gaiotto states with fundamental multiplets through Degenerate DAHA,* JHEP **1409**, 028 (2014) [arXiv:1405.3141 [hep-th]].

[20] P. Argyres and M. Douglas, *New phenomena in SU(3) supersymmetric gauge theory*, Nucl. Phys. **B 448** (1995) 93 [hep-th/95055062].

[21] P. Argyres, M. Plesser, N. Seiberg and E. Witten, *New N=2 superconformal field theories in four dimensions*, Nucl. Phys. **B 461** (1996) 71 [hep-th/9511154].

[22] L. F. Alday, D. Gaiotto and Y. Tachikawa, *Liouville Correlation Functions from Four-dimensional Gauge Theories*, Lett. Math. Phys. **91**, 167 (2010) [arXiv:0906.3219].

[23] D. Gaiotto, G. W. Moore, and A. Neitzke, *Wall-crossing, Hitchin System, and the WKB Approximation* [arXiv:0907.3987].

[24] G. Bonelli and A. Tanzini, *Hitchin systems, N=2 gauge theories and W-gravity*, Physl. Lett. **B691** (2010) 111 [arXiv:0909.4031].

[25] D. Nanopoulos and D. Xie, *Hichin equation, singularity, and N=2 superconformal field theories*, JHEP **1003** (2010) 043 [arXiv:0911.1990].